

28 February 2025





Getting to know Blue Cone Monochromacy

Dear ophthalmologists, low-vision specialists, optometrists, geneticists, pediatricians, other medical professionals, and friends of BCM:

28 February 2025 marks the international Rare Disease Day coordinated by European Organisation for Rare Diseases (EURORDIS) and National Organization for Rare Disorders (NORD).

Today patients, families, and volunteers of BCM Families Foundation reached you with this flyer because we would like to tell you about Blue Cone Monochromacy, also known as BCM, a rare genetic eye condition affecting 1 in 100,000 people.

BCM Families Foundation is a non-profit 501(c)(3) organization, founded and registered in the US by patients affected by BCM and their relatives. Our mission is to cure Blue Cone Monochromacy.

BCM at a glance

Blue Cone Monochromacy is a rare retinal disorder and is inherited in an X-linked manner. Affected males commonly show symptoms soon after birth and BCM is thought to be a relatively stable disease, affecting an individual for his whole life. Female carriers rarely show symptoms, and when they do, they are very mild.

Symptoms of BCM

Poor visual acuity ranging from 20/60 to 20/200 or greater, in some cases legal blindness.

Poor or no color discrimination, Patients with BCM have functioning rod photoreceptors and blue cone photoreceptors. However, they lack function of their red (L) and green (M) cone photoreceptors.

Hemeralopia, Patients with BCM have severe difficulty coping with glare because they only have blue cone function at higher light levels. The severity of the symptoms, commonly known as photophobia, varies by the integrity of remaining blue cone cells.


Nystagmus: Nystagmus is usually present from about 3-6 months of age. It is usually, but not always, pendular and may decrease with age.


Myopia or high myopia

X-linked inheritance family pedigree. Mutations causing BCM are located in the X chromosome in the following genes and supporting components: **OPNILW, OPNIMW** and the upstream **LCR** (Locus Control Region). Deletion in the LCR or a point mutation C203R on a single L-M opsin gene are most common forms.

So if you see a 6 month old boy with nystagmus and the parents speak about another male in their family (on the mother's side) with similar eye problems, please consider a diagnosis of BCM.

X-Linked recessive Pedigree





For confirmatory diagnosis, please conduct genetic testing at:

University of Tuebingen Molecular Genetics Laboratory - Germany Contact: Susanne Kohl, Msc, PhD Email: susanne.kohl@uni-tuebingen.de No-Cost DNA Test Program for Blue Cone Monochromacy https://www.blueconemonochromacy.org/no-cost-dna-test/

Molecular Vision Laboratory

1920 NE Stucki Ave, Suite 150 Hillsboro, OR 97006 Contact: John Chiang, PhD, FACMG Email: jchiang@mvisionlab.com Web: www.molecularvisionlab.com **Radboud Medical Genetics Institute** Nijmegen The Netherland Email: info@gdnm.nl Web: https://order.radboudumc.nl

John and Marcia Carver Nonprofit Genetic Testing Laboratory University of Iowa 375 Newton Rd. 4111 MERF Iowa City, IA 52242 Email: carverlab@uiowa.edu Web: https://www.carverlab.org

Participate in the research supported by BCM Families Foundation at:

University of Pennsylvania - Dr. A. Cideciyan and Dr. T.S. Aleman https://www.med.upenn.edu/chrd/

BCMFF Collaboration

Medical College of Wisconsin - Dr. J. Carroll Web:https://www.mcw.edu/departments/ophthalmology-eye-institute/research/advanced-ocular-imaging-program

For more information about the disease, we invite you to visit the Online Mendelian Inheritance in Man (OMIM) page for BCM (http://omim.org/entry/303700) and our website (www.bcmfamilies.org)

BCM Families Foundation PO Box 7711 Jupiter, FL 33468-7711 USA Email: info@BCMFamilies.org Web: www.bcmfamilies.org



Forum: bcmfamilies.ning.com Facebook: BCM Families Foundation

